arXiv:1512.05682v1 [math.CO] 17 Dec 2015

k-connected degree sequences

Jonathan McLaughlin

Abstract. Necessary and sufficient conditions for a sequence of positive integers to be the degree sequence of a k-connected simple graph are detailed. Conditions are also given under which such a sequence is necessarily k-connected.

Mathematics Subject Classification (2010). Primary 05C40.

Keywords. k-connected graph, degree sequence.

1. Introduction

Necessary and sufficient conditions for a sequence of non-negative integers to be connected i.e. the degree sequence of some finite simple connected graph, are implicit in Hakimi [4] and have been stated explicitly by the author in [5] and expounded on in [6]. This note builds upon these conditions of Hakimi and details necessary and sufficient conditions for a sequence of non-negative integers to be k-connected. The note concludes with necessary and sufficient conditions for a sequence of non-negative integers to be necessarily k-connected i.e. the sequence can only be realised as a k-connected graph.

2. Preliminaries

Let $G = (V_G, E_G)$ be a graph where V_G denotes the vertex set of G and $E_G \subseteq [V_G]^2$ denotes the edge set of G (given that $[V_G]^2$ is the set of all 2-element subsets of V_G). An edge $\{a, b\}$ is denoted ab. A graph is finite when $|V_G| < \infty$ and $|E_G| < \infty$, where $|X|$ denotes the cardinality of the set X. The union of graphs G and H is the graph $G \cup H = (V_G \cup V_H, E_G \cup E_H)$ and $G \cup ab$ is understood to be the graph $(V_G, E_G) \cup (\{a, b\}, \{ab\})$. A graph is simple if it contains no loops (i.e. $aa \notin E_G$) or parallel/multiple edges (i.e. $\{ab, ab\} \not\subseteq E_G$). The *degree* of a vertex v in a graph G, denoted $deg(v)$, is the number of edges in G which contain v. A graph where all vertices have degree k is called a *k-regular* graph. A *path* is a graph with n vertices in which two vertices, known as the *endpoints*, have degree 1 and $n-2$ vertices

have degree 2. A graph is *connected* if there exists at least one path between every pair of vertices in the graph. Paths P_1 and P_2, both with endpoints a and b, are *internally disjoint* if $P_1 \cap P_2 = (\{a, b\}, \{\})$. A graph G is *k-connected* when there exists at least k internally disjoint paths in G between any two vertices in G. This characterisation of a graph being k-connected is synonymous with Menger's Theorem. K_n denotes the *complete graph* on n vertices. The *complement* of a simple graph G is the simple graph $\overline{G}$ with vertex set V_G and edge set the pairs of vertices in V_G which are not contained in E_G. All basic graph theory definitions can be found in standard texts such as [1], [2] or [3].

3. Degree sequences

A finite sequence $s = \{s_1, ..., s_n\}$ of non-negative integers is called *graphic* if there exists a finite simple graph with vertex set $\{v_1, ..., v_n\}$ such that v_i has degree s_i for all $i = 1, ..., n$. Given a graph G then the degree sequence $d(G)$ is the monotonic non-increasing sequence of degrees of the vertices in V_G. This means that every graphical sequence s is equal to the degree sequence $d(G)$ of some graph G (subject to possible rearrangement of the terms in s).

Definition 3.1. A finite sequence $s = \{s_1, ..., s_n\}$ of positive integers is called *k-connected* if there exists a finite simple k-connected graph with vertex set $\{v_1, ..., v_n\}$ such that $deg(v_i) = s_i$ for all $i = 1, ..., n$.

A finite sequence $s = \{s_1, ..., s_n\}$ of positive integers is called *necessarily k-connected* if s can only be realisable as a simple k-connected graph.

Given a sequence of positive integers $s = \{s_1, ..., s_n\}$ then define the *associated pair of* s, denoted $(\varphi(s), \epsilon(s))$, to be the pair $(n, \frac{1}{2}\sum_{i=1}^{n} s_i)$. Where no ambiguity can arise, $(\varphi(s), \epsilon(s))$ is simply denoted (φ, ϵ).

4. Results

Theorem 4.1. *Given a sequence $s = \{s_1, ..., s_n\}$ of positive integers, with the associated pair (φ, ϵ), such that $s_i \geq s_{i+1}$ for $i = 1, ..., n-1$ then s is k-connected if and only if*

- $\epsilon \in \mathbb{N}$,
- $s_1 \leq \varphi - 1$ *and* $s_n \geq k$,
- $\frac{k\varphi}{2} \leq \epsilon \leq \binom{\varphi}{2}$.

Proof. ($\Rightarrow$) Clearly $\epsilon \in \mathbb{N}$ is a necessary condition for any sequence s to be realisable as half the sum of the degrees in any graph is the number of edges in that graph which must be a natural number. The necessity of the condition $s_1 \leq \varphi - 1$ follows from the definition of a simple graph and the need for $s_n \geq k$ is due to the fact that every vertex in a k-connected graph has degree

at least k.

The necessity of the condition $\epsilon \geq \frac{k\varphi}{2}$ follows from the observation that the minimum possible ϵ of any k-connected sequence is $\epsilon = \frac{k\varphi}{2}$ which is contained in the associated pair of $s = \underbrace{\{k, ..., k\}}_{\varphi}$, the degree sequence of a k-regular graph. Note that ϵ must be even, by definition, and this occurs whenever φ and k are not *both* odd (as there cannot exist a k-regular graph, where k is odd and the number of vertices is also odd). If both φ and k are odd, then the minimum possible ϵ of any k-connected sequence is $\epsilon = \frac{k\varphi+1}{2}$ which is contained in the associated pair of $s = \underbrace{\{k+1, k, ..., k\}}_{\varphi}$. To show the necessity of the condition $\epsilon \leq \binom{\varphi}{2}$, the parity of φ is irrelevant. When maximising ϵ it follows that the maximum possible ϵ of any k-connected sequence is $\epsilon = \binom{\varphi}{2}$ which is contained in the associated pair of $s = \underbrace{\{\varphi - 1, ..., \varphi - 1\}}_{\varphi}$, the degree sequence of the complete graph K_φ.

It remains to show that, for a fixed φ, all sequences with $\frac{k\varphi}{2} < \epsilon < \binom{\varphi}{2}$ are realisable. Suppose that φ and k are not both odd and $k < \varphi - 1$. To ensure that ϵ remains even, two terms in $\{k, ..., k\}$ must be incremented by one to give $\{k+1, k+1, k, ..., k\}$. Given that $\{k, ..., k\}$ is realisable as a k-regular graph G then $\{k+1, k+1, k, ..., k\}$ is realisable as $G \cup ab$ where $ab \in \overline{G}$. This incrementing of two terms in a graphic sequence can be continued until the sequence with $\epsilon = \binom{\varphi}{2}$ is reached and this process is summarised in FIGURE 1.

$s = \{s_1, \ldots, s_n\}$	ϵ	$\lvert E_G \rvert$
$\{k, \ldots, k\}$	$\frac{k\varphi}{2}$	$\frac{kn}{2}$
$\{k+1, k+1, k, \ldots, k\}$	$\frac{k\varphi+2}{2}$	$\frac{kn}{2} + 1$
$\vdots$	$\vdots$	$\vdots$
$\{n-1, \ldots, n-1\}$	$\binom{\varphi}{2}$	$\binom{n}{2}$

FIGURE 1. k-connected degree sequences when φ and k are not both odd.

A similar argument exists when φ and k are both odd, with $k < \varphi - 1$, and this argument is summarised in FIGURE 2.

($\Leftarrow$) Suppose that $s = \{s_1, ..., s_n\}$ is k-connected. This means that s is the degree sequence of a k-connected graph G, hence $\sum_{i=1}^{n} deg(v_i) = 2|V_G|$ and

$s = \{s_1, \ldots, s_n\}$	ϵ	$\lvert E_G \rvert$
$\{k+1, k, \ldots, k\}$	$\frac{k\varphi+1}{2}$	$\frac{kn+1}{2}$
$\{k+2, k+1, k, \ldots, k\},\ \{k+1, k+1, k+1, k, \ldots, k\}$	$\frac{(k\varphi+1)+2}{2}$	$\frac{kn+1}{2}+1$
$\vdots$	$\vdots$	$\vdots$
$\{n-1, \ldots, n-1\}$	$\binom{\varphi}{2}$	$\binom{n}{2}$

FIGURE 2. k-connected degree sequences when φ and k are both odd.

so $\epsilon \in \mathbb{N}$. As G is k-connected then $deg(v_i) \geq k$ for all $i = 1, ..., n$ hence if G is a minimally k-connected graph on n vertices then $d(G) = \{k, ..., k\}$ with $|E_G| = \frac{kn}{2}$ if n and k are not both odd or $d(G) = \{k+1, k, ..., k\}$ with $|E_G| = \frac{kn+1}{2}$ if n and k are both odd, hence $s_n \geq k$ and $\epsilon \geq \frac{k\varphi}{2}$. As G is simple then $deg(v_i) \leq n-1$ for all $i = 1, ..., n$ and the maximal simple (k-connected) graph on n vertices is the complete graph K_n which has the degree sequence $\{n-1, ..., n-1\}$ and $|E_{K_n}| = \binom{n}{2}$, hence $s_1 \leq n-1$ and $\epsilon \leq \binom{\varphi}{2}$. To show that k-connected graphs exist for each $|E_G| \in \mathbb{N}$ such that $\frac{kn}{2} < |E_G| < \binom{n}{2}$ refer to the argument showing the existence of graphic sequences with $\frac{k\varphi}{2} < \epsilon < \binom{\varphi}{2}$ detailed above, along with FIGURES 1 and 2. □

Theorem 4.2. *Given a sequence $s = \{s_1, ..., s_n\}$ of positive integers, with the associated pair (φ, ϵ), such that $s_i \geq s_{i+1}$ for $i = 1, ..., n-1$ then s is necessarily k-connected if and only if s is k-connected and $\epsilon > \binom{\varphi-2}{2} + 2k - 1$.*

Proof. ($\Rightarrow$) Clearly it is necessary for s to be k-connected if it is to be necessarily k-connected. It is required to show that it is necessary for $\epsilon > \binom{\varphi-2}{2} + 2k - 1$. Consider a sequence $s = \{s_1, ..., s_n\}$ such that $\epsilon = \binom{\varphi-2}{2} + 2k - 1$.

Observe that one such sequence is $s' = \{\underbrace{n-1, ..., n-1}_{k-1}, \underbrace{n-3, ..., n-3}_{n-k-1}, \underbrace{k, k}_{2}\}$ where $\varphi(s') = k-1+n-k-1+2 = n$ and $\epsilon(s') = \frac{(k-1)(n-1)+(n-k-1)(n-3)+2k}{2}$ $= \frac{(n-k-1+k-1)(n-3)+2(k+k-1)}{2} = \frac{(n-2)(n-3)+2(2k-1)}{2} = \binom{n-2}{2} + 2k - 1$.

Observe that $s' = d(G_1)$, see FIGURE 3, where $G_1 = H_1 \cup H_2$ with $H_1 \simeq K_{k+1}$, $H_2 \simeq K_{n-2}$ and $H_1 \cap H_2 \simeq K_{k-1}$. Note that as $G_1 \setminus (H_1 \cap H_2)$ is disconnected and that $H_1 \cap H_2 \simeq K_{k-1}$ (i.e. $|V_{H_1 \cap H_2}| = k-1$) then G_1 is $(k-1)$-connected.

However, $s' = \{\underbrace{n-1, ..., n-1}_{k-1}, \underbrace{n-3, ..., n-3}_{n-k-1}, \underbrace{k, k}_{2}\}$ is, in fact, k-connected as s' is also the degree sequence of G_2, see FIGURE 3 (noting that $v_i v_j \in E_{G_1}$ but $v_i v_j \notin E_{G_2}$). Therefore, it is required that $\epsilon > \binom{n-2}{2} + 2k - 1$ if s is to



FIGURE 3. For $k \geq 1$ and $s' = \{n-1, ..., n-1, n-3, ..., n-3, k, k\} = d(G_1) = d(G_2)$.

be necessarily k-connected.

Note that FIGURES 4, 5 and 6 are, respectively, the $k = 1, 2$ and 3 versions of FIGURE 3.

FIGURE 4. $k = 1$ and $s' = \{n-3, ..., n-3, 1, 1\} = d(G_1) = d(G_2)$.

FIGURE 5. $k = 2$ and $s' = \{n-1, n-3, ..., n-3, 2, 2\} = d(G_1) = d(G_2)$.

($\Leftarrow$) It is now required to show that if s is k-connected and $\epsilon > \binom{n-2}{2} + 2k - 1$ then s is necessarily k-connected. To show this it is required to show that the maximum number of edges in a graph with n vertices which is not k-connected is $\binom{n-2}{2} + 2k - 1$. The graph G_1 in FIGURE 3 shows that such a graph exists, so it remains to show that a graph with $\epsilon = \binom{n-2}{2} + 2k - 1$ is maximally $(k-1)$-connected i.e. adding one edge will always result in a k-connected graph.



FIGURE 6. $k = 3$ and $s' = \{n-1, n-1, n-3, ..., n-3, 3, 3\} = d(G_1) = d(G_2)$.

Observe that any maximally $(k-1)$-connected graph G on n vertices will necessarily contain a cut set C containing $k-1$ vertices. This means that $G \setminus C$ is disconnected. To maximise the number of edges in G it is clear that $G \setminus C$ contains two connected components i.e. $G = H_1 \cup H_2$ where $V_{H_1} \cap V_{H_2} = C$ with $H_1 \simeq K_{a+|C|}$, $H_2 \simeq K_{b+|C|}$ and $H_1 \cap H_2 \simeq K_{|C|}$ (noting that $a+b = n-|C|$). So, the task of maximising $|E_{H_1 \setminus C}| + |E_{H_2 \setminus C}|$ is equivalent to minimising the number of edges in a complete bipartite graph $K_{a,b}$ as $K_n \setminus (E_{H_1 \cup H_2}) \simeq K_{a,b}$.

Let $a + b = n - |C|$, with $a \leq b$, then $|E_{K_{a,b}}| = ab$ where $a, b \in \{1, ..., n - |C| - 1\}$. Note that $a > 0$ as $G \setminus C$ is disconnected i.e. $K_a \neq K_0 = (\varnothing, \varnothing)$. It is straightforward to show that ab attains its maximum at $a = b = \frac{n-|C|}{2}$, when n is even, and at $a = \lfloor \frac{n-|C|}{2} \rfloor, b = \lceil \frac{n-|C|}{2} \rceil$ when n is odd. It follows that ab is minimised when $a = 1$ and $b = n - |C| - 1$. However, observe that $a > 1$ as $a = 1$ implies that $H_1 \simeq K_{|C|+1}$ which means that $d(G)$ contains a term equal to $|C| = k - 1$, but this contradicts the $s_n \geq k$ condition. Hence $|E_{K_{a,b}}|$, with $a + b = n - |C|$, is minimised when $a = 2$ and $b = n - |C| - 2$ and so any maximally $(k-1)$-connected graph on n vertices is isomorphic to $H_1 \cup H_2$ where $H_1 \simeq K_{k+1}, H_2 \simeq K_{n-k-1}$ and $H_1 \cap H_2 \simeq K_{k-1}$, see G_1 in FIGURE 3. Notice that the union of G_1 and any edge in $\overline{G_1}$ results in a k-connected graph. □

Corollary 4.3. *All simple graphs with n vertices and at least $\frac{n^2-5n+6+4k}{2}$ edges are k-connected.*

Proof. As shown in Theorem 4.2, a maximally $(k-1)$-connected graph with n vertices is isomorphic to the union of $H_1 \simeq K_{k+1}$ and $H_2 \simeq K_{n-2}$ where $H_1 \cap H_2 \simeq K_{k-1}$ and such a graph has $|E_{K_{k+1}}| + |E_{K_{n-2}}| - |E_{K_{k-1}}| = \binom{n-2}{2} + \binom{k+1}{2} - \binom{k-1}{2} = \binom{n-2}{2} + \frac{(k+1)k}{2} - \frac{(k-1)(k-2)}{2} = \binom{n-2}{2} + \frac{4k-2}{2}$ edges. Hence, any simple graph with n vertices and at least $\left(\binom{n-2}{2} + 2k - 1\right) + 1 = \frac{(n-2)(n-3)+4k}{2}$ edges is k-connected. □

Note that for all $n, k \in \mathbb{N}$, $n^2 - 5n$ is even and $n^2 - 5n + 6 + 4k > 0$, hence $\frac{n^2-5n+6+4k}{2} \in \mathbb{N}$.

References

[1] Bondy, J. A. and Murty, U. S. R., *Graph theory,* Graduate Texts in Mathematics **244** (2008), Springer.

[2] Diestel, R., *Graph theory,* Graduate Texts in Mathematics **173** (2000), Springer-Verlag.

[3] Gould, R., *Graph theory,* (1988), The Benjamin/Cummings Publishing Co. Inc.

[4] Hakimi, S. L., *On realizability of a set of integers as degrees of the vertices of a linear graph. I,* J. Soc. Indust. Appl. Math. **10** (1962), 496–506.

[5] McLaughlin, J., *On connected degree sequences,* arXiv:1511.09321 (2015).

[6] McLaughlin, J., *On connected simple graphs and their degree sequences,* arXiv:1512.01377 (2015).

Jonathan McLaughlin
Department of Mathematics,
St. Patrick's College,
Dublin City University,
Dublin 9,
Ireland
e-mail: jonny_mclaughlin@hotmail.com